




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 207/40, 409/12, 409/14, A01N 43/36	A1	(11) International Publication Number: WO 00/09482 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/EP99/05837 (22) International Filing Date: 10 August 1999 (10.08.99) (30) Priority Data: 9817548.2 12 August 1998 (12.08.98) GB (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (72) Inventors; and (75) Inventors/Applicants (for US only): EBERLE, Martin [CH/CH]; Asterhagstrasse 22, CH-4103 Bottmingen (CH). WALTER, Harald [DE/CH]; Chilchmattstrasse 12b, CH-4118 Rodersdorf (CH). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: TRIFLUORO METHYLPYRROL CARBOXAMIDES**(57) Abstract**

The invention relates to novel pesticidal trifluoromethylpyrrolcarboxamides of formula (I), wherein R₁ is hydrogen, halogen, C₁-4haloalkyl, or C₁-4alkyl, R₂ is C₁-4alkyl, C₁-4haloalkyl, C₁-4alkoxy-C₁-4alkyl, cyano, C₁-4alkylsulfonyl, phenylsulfonyl, di(C₁-4alkyl)aminosulfonyl, C₁-6alkylcarbonyl, benzoyl, or substituted phenylsulfonyl or benzoyl, and A is a group (1), (2) or (3) wherein R₃ is C₁-6alkyl, C₁-6haloalkyl, C₂-6alkenyl, C₂-6haloalkenyl, C₂-6alkinyl, C₁-6alkoxy, C₁-6haloalkoxy, C₂-6alkenyloxy, C₂-6haloalkenyloxy, C₂-6alkinyloxy, C₃-7cycloalkyl, C₁-4alkyl-C₃-7cycloalkyl, C₄-7cycloalkenyl, C₁-4alkyl-C₄-7cycloalkenyl, C₃-7cycloalkyloxy, C₁-4alkyl-C₃-7cycloalkyloxy, C₅-7cycloalkenyloxy, C₁-4alkyl-C₅-7cycloalkenyloxy, phenyl, naphthyl, phenoxy, naphthyloxy, or substituted phenyl or phenoxy wherein the substituents are one to three groups independently selected from halogen, C₁-4alkyl, C₁-4alkoxy, C₁-4alkylthio, cyano, C₁-4alkoxycarbonyl, C₁-4alkylcarbonyl, C₁-4haloalkyl, C₁-4haloalkoxy, methylenedioxy or difluoromethylenedioxy, or phenyl; R₄ is hydrogen, halogen C₁-4alkyl, C₁-4haloalkyl, C₁-4alkoxy or C₁-4haloalkoxy; and R₅, R₆ and R₇ independently of each other are C₁-6alkyl, C₃-7cycloalkyl or C₃-7cycloalkyl-C₁-4alkyl. The novel compounds have plant-protecting properties and are suitable for protecting plants against infestation by phytopathogenic microorganisms.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

TRIFLUOROMETHYLPYRROLCARBOXAMIDES

The present invention relates to novel trifluoromethylpyrrolicarboxamides which have microbicidal activity, in particular fungicidal activity. The invention also relates to the preparation of these substances, to agrochemical compositions which comprise at least one of the novel compounds as active ingredient, to the preparation of the compositions mentioned and to the use of the active ingredients or compositions in agriculture and horticulture for controlling or preventing infestation of plants by phytopathogenic microorganisms, preferably fungi.

The trifluoromethylpyrrolicarboxamides of the present invention have the general formula I




wherein

R₁ is hydrogen, halogen, C₁₋₄haloalkyl or C₁₋₄alkyl,

R₂ is C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy-C₁₋₄alkyl, cyano, C₁₋₄alkylsulfonyl, phenylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, C₁₋₆alkylcarbonyl, benzoyl, or substituted phenylsulfonyl or benzoyl, and

A is a group



wherein

R₃ is C₁₋₆alkyl, C₁₋₆haloalkyl, C₂₋₆alkenyl, C₂₋₆haloalkenyl, C₂₋₆alkinyl, C₁₋₆alkoxy, C₁₋₆haloalkoxy, C₂₋₆alkenyloxy, C₂₋₆haloalkenyloxy, C₂₋₆alkinyloxy, C₃₋₇cycloalkyl, C₁₋₄alkyl-C₃₋₇cycloalkyl, C₄₋₇cycloalkenyl, C₁₋₄alkyl-C₄₋₇cycloalkenyl, C₃₋₇cycloalkyloxy, C₁₋₄alkyl-C₃₋₇cycloalkyloxy, C₅₋₇cycloalkenyloxy, C₁₋₄alkyl-C₅₋₇cycloalkenyloxy, phenyl, naphthyl, phenoxy, naphthyloxy, or substituted phenyl or phenoxy wherein the substituents are one to three groups independently selected from halogen, C₁₋₄alkyl,

C₁₋₄alkoxy, C₁₋₄alkylthio, cyano, C₁₋₄alkoxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄haloalkyl, C₁₋₄haloalkoxy, methylenedioxy or difluoromethylenedioxy, or phenyl;

R₄ is hydrogen, halogen, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy or C₁₋₄haloalkoxy ; and

R₅, R₆ and R₇ independently of each other are C₁₋₆alkyl, C₃₋₇cycloalkyl or

C₃₋₇cycloalkyl-C₁₋₄alkyl.

Nicotinic acid amides with fungicidal properties useful in agricultural practice are known from EP-A-0545099. The disclosed compounds for practical purposes do not always satisfy the needs of the modern agriculture.

Surprisingly , it has now been found that the compounds of formula I exhibit improved biological properties which render them more suitable for the practical use in agriculture and horticulture.

Where asymmetrical carbon atoms are present in the compounds of formula I, these compounds are in optically active form. The invention relates to the pure isomers, such as enantiomers and diastereomers, as well as to all possible mixtures of isomers, e.g. mixtures of diastereomers, racemates or mixtures of racemates.

Within the present specification alkyl denotes methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, iso-pentyl, sec-pentyl, tert-pentyl, and neo-pentyl. Non-branched alkyl is preferred. Alkyl as part of other radicals such as alkoxy, alkylthio, haloalkyl, alkylcycloalkyl, alkylcycloalkoxy, alkylcarbonyl, alkylsulfonyl, alkylamino, etc. is understood in an analogous way. Halogen will be understood generally as meaning fluoro, chloro, bromo or iodo. Fluoro, chloro or bromo are preferred meanings. Halogen as part of other radicals such as haloalkyl, haloalkoxy, haloalkenyl, haloalkenyloxy or halophenyl, etc. is understood in an analogous way. Substituents of the substituted phenylsulfonyl and benzoyl may be present mono- to five times and are preferably independently selected from halogen, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylthio, cyano, C₁₋₄alkoxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄haloalkyl, C₁₋₄haloalkoxy, methylenedioxy or difluoromethylenedioxy, or phenyl. Where phenyl groups as parts of phenyl, phenylsulfonyl, phenoxy and benzoyl are substituted such radicals for example designate 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,4-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 3,5-difluorophenyl,


2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-trifluoromethoxyphenyl, 3-trifluoromethoxyphenyl, 4-trifluoromethoxyphenyl, 4-methylphenyl, 4-ethylphenyl, 4-isopropylphenyl, 4-tert.butylphenyl, 3,4-dimethylphenyl, 2,4,6-trimethylphenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-biphenyl, 3-biphenyl, and the like.

A special subgroup of active ingredients is represented by those compounds of formula I wherein

R₁ is hydrogen or C₁₋₄alkyl,

R₂ is C₁₋₄alkyl, and

A is a group



wherein

R₃ is C₁₋₆alkyl, C₁₋₆haloalkyl, C₂₋₆alkenyl, C₂₋₆haloalkenyl, C₂₋₆alkinyl, C₁₋₆alkoxy, C₁₋₆haloalkoxy, C₂₋₆alkenyloxy, C₂₋₆haloalkenyloxy, C₂₋₆alkinyloxy, C₃₋₇cycloalkyl, C₁₋₄alkyl-C₃₋₇cycloalkyl, C₄₋₇cycloalkenyl, C₁₋₄alkyl-C₄₋₇cycloalkenyl, C₃₋₇cycloalkyloxy, C₁₋₄alkyl-C₃₋₇cycloalkyloxy, C₅₋₇cycloalkenyloxy, C₁₋₄alkyl-C₅₋₇cycloalkenyloxy, phenyl, naphthyl, phenyl substituted by one to three groups independently selected from halogen, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylthio, cyano, C₁₋₄alkoxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄haloalkyl, C₁₋₄haloalkoxy, methylenedioxy or difluoromethylenedioxy, or phenyl;

R₄ is hydrogen, halogen or C₁₋₄alkyl ;and

R₅, R₆ and R₇ independently of each other are C₁₋₆alkyl, C₃₋₇cycloalkyl or C₃₋₇cycloalkyl-C₁₋₄alkyl.

Within the group of compounds of formula I those compounds are preferred wherein:

- R₁ is hydrogen or methyl, or
- R₂ is methyl, or
- R₃ is phenyl or phenyl substituted with halogen.


Another preferred subgroup is constituted by the compounds of formula I wherein R₁ is hydrogen or methyl, R₂ is methyl, and R₃ is phenyl or phenyl substituted with halogen.


Preferred individual compounds are:

N-(2-biphenyl)-1-methyl-4-trifluoromethylpyrrol-3-carboxamide,
 N-(4'-chloro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide,
 N-(4'-fluoro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide,
 N-[2-(4-fluorophenyl)-3-thienyl]-4-trifluoromethylpyrrol-3-carboxamide,
 N-[2-(4-chlorophenyl)-3-thienyl]-4-trifluoromethylpyrrol-3-carboxamide,
 N-(3',4'-difluoro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide,
 N-(3'-trifluoromethyl-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide, and
 N-(4'-chloro-3'-fluoro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide.

The compounds according to formula I may be prepared according to the following reaction scheme 1.

Scheme 1:






wherein A, R₁ and R₂ are as defined for formula I, X and Y are leaving groups, alkyl designates a lower alkyl moiety, and aryl stands for phenyl or tolyl.

The leaving groups X and Y preferably designate halogen atoms, but may also stand for moieties of mixed anhydrides, e.g. -O-CO-alkyl, -O-PO(alkyl)₂, -O-C(=N-alkyl), triazolyl, and the like.

Alternatively the introduction of the radicals R₂ and A may be a different order according to scheme 2.

Scheme 2:



wherein A, R₁, R₂, X and Y are as defined for scheme 1

In special cases of substitution patterns it may be favourable to utilize a protective group at the 1-position of the pyrrol ring in order to avoid side reactions during the amidation reaction at the carbonyl function in 3-position. Suitable protective groups are benzyl and the like. Also introduction and cleaving of the protective group is done by common benzylation and hydrolysis reactions.

The new carboxamides of formula I are preferably obtained by reacting an activated carboxylic acid of formula II with an aromatic amine. Depending on the leaving group Y the reaction is run in the presence of an acid binding agent. Inert solvents are preferably used, but also aqueous solvent mixtures, referred to as Schotten-Baumann conditions, may be employed.

Activation is achieved by reacting the pyrrol carboxylic acid III with an activating agent such as acid halide, such as thionyl chloride or oxalyl chloride, to give the corresponding acyl chloride (Y=chlorine) or by any other general coupling activation method known in the literature.

The carboxylic acids of formula III may be prepared from the corresponding esters by hydrolysis using standard hydrolysis conditions.

The N-alkylation of pyrrol carboxylic esters of formula V to give the N-alkylated pyrrol carboxylic esters of formula IV may be achieved by reacting a pyrrol carboxylic ester of formula V with an alkylating agent, such as a dialkylsulfate, an alkyl halogenide or an alkylsulfonate using standard conditions. The pyrrol carboxylic esters of formula V may typically be prepared according to the van Leusen protocol, i.e. the reaction of an alkyl acrylate with an aryl sulfonyl isocyanide in the presence of a base. Typical bases employed in scheme 1 include metal hydrides, metal alcoholates and metal hydroxides. Any solvent which is inert under the reaction conditions may be used. Representative solvents include ethers (diethylether, dimethoxyethane, THF and the like), polar aprotic solvents, such as DMSO, DMF, DMA, or protic solvents, such as alcohols. Also mixtures of such solvents may be used. In many cases phase transfer reaction conditions are applicable which naturally require a two-phase reaction system. Preferably the entire reaction sequence of scheme 1 is conducted as a single-vessel-reaction.

Surprisingly, it has now been found that the novel compounds of formula I have, for practical purposes, a very advantageous spectrum of activities for protecting plants against diseases that are caused by fungi as well as by bacteria and viruses.

The compounds of formula I can be used in the agricultural sector and related fields of use as active ingredients for controlling plant pests. The novel compounds are distinguished by excellent activity at low rates of application, by being well tolerated by plants and by being environmentally safe. They have very useful curative, preventive and systemic properties and are used for protecting numerous cultivated plants. The compounds of formula I can be used to inhibit or destroy the pests that occur on plants or parts of plants (fruit, blossoms, leaves, stems, tubers, roots) of different crops of useful plants, while at the same time protecting also those parts of the plants that grow later e.g. from phytopathogenic micro-organisms.

It is also possible to use compounds of formula I as dressing agents for the treatment of plant propagation material, in particular of seeds (fruit, tubers, grains) and plant cuttings (e.g. rice), for the protection against fungal infections as well as against phytopathogenic fungi occurring in the soil.

The compounds I are, for example, effective against the phytopathogenic fungi of the following classes: Fungi imperfecti (e.g. Botrytis, Pyricularia, Helminthosporium, Fusarium, Septoria, Cercospora and Alternaria) and Basidiomycetes (e.g. Rhizoctonia, Hemileia, Puccinia). Additionally, they are also effective against the Ascomycetes classes (e.g. Venturia and Erysiphe, Podosphaera, Monilinia, Uncinula) and of the Oomycetes classes (e.g. Phytophthora, Pythium, Plasmopara). Outstanding activity has been observed against powdery mildew (Erysiphe spp.). Furthermore, the novel compounds of formula I are effective against phytopathogenic bacteria and viruses (e.g. against Xanthomonas spp, Pseudomonas spp, Erwinia amylovora as well as against the tobacco mosaic virus).

Within the scope of present invention, target crops to be protected typically comprise the following species of plants: cereal (wheat, barley, rye, oat, rice, maize, sorghum and related species); beet (sugar beet and fodder beet); pomes, drupes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries); leguminous

plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (pumpkins, cucumbers, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, mandarins); vegetables (spinache, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocado, cinnamomum, camphor) or plants such as tobacco, nuts, coffee, eggplants, sugar cane, tea, pepper, vines, hops, bananas and natural rubber plants, as well as ornamentals.

The compounds of formula I are used in unmodified form or, preferably, together with the adjuvants conventionally employed in the art of formulation. To this end they are conveniently formulated in known manner to emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations e.g. in polymeric substances. As with the type of the compositions, the methods of application, such as spraying, atomising, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances. The compositions may also contain further adjuvants such as stabilisers, antifoams, viscosity regulators, binders or tackifiers as well as fertilisers, micronutrient donors or other formulations for obtaining special effects.

Suitable carriers and adjuvants can be solid or liquid and are substances useful in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilisers. Such carriers are for example described in WO 97/33890.

The compounds of formula I are normally used in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession with further compounds. These further compounds can be e.g. fertilisers or micronutrient donors or other preparations which influence the growth of plants. They can also be selective herbicides as well as insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures of several of these preparations, if desired together with further carriers, surfactants or application promoting adjuvants customarily employed in the art of formulation.

The compounds of formula I can be mixed with other fungicides, resulting in some cases in unexpected synergistic activities.

Mixing components which are particularly preferred are azoles such as azaconazole, bitertanol, propiconazole, difenoconazole, diniconazole, cyproconazole, epoxiconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imazalil, imibenconazole, ipconazole, tebuconazole, tetraconazole, fenbuconazole, metconazole, myclobutanil, perfurazoate, penconazole, bromuconazole, pyrifenox, prochloraz, triadimefon, triadimenol, triflumizole or triticonazole; pyrimidinyl carbinols such as ancymidol, fenarimol or nuarimol; 2-amino-pyrimidine such as bupirimate, dimethirimol or ethirimol; morpholines such as dodemorph, fenpropidin, fenpropimorph, spiroxamin or tridemorph; anilinopyrimidines such as cyprodinil, pyrimethanil or mepanipyrim; pyrroles such as fenciclonil or fludioxonil; phenylamides such as benalaxyl, furalaxyl, metalaxyl, R-metalaxyl, ofurace or oxadixyl; benzimidazoles such as benomyl, carbendazim, debacarb, fuberidazole or thiabendazole; dicarboximides such as chlozolate, dichlozoline, iprodione, myclozoline, procymidone or vinclozolin; carboxamides such as carboxin, fenfuram, flutolanil, mepronil, oxycarboxin or thifluzamide; guanidines such as guazatine, dodine or iminoctadine; strobilurines such as azoxystrobin, kresoxim-methyl, metominostrobin, SSF-129, methyl 2-[(2-trifluoromethyl)-pyrid-6-yloxymethyl]-3-methoxyacrylate or 2-[α [(α -methyl-3-trifluoromethyl-benzyl)imino]-oxy]-o-tolyl]-glyoxylic acid-methylester-O-methyloxime (trifloxystrobin); dithiocarbamates such as ferbam, mancozeb, maneb, metiram, propineb, thiram, zineb or ziram; N-halomethylthio-dicarboximides such as captafol, captan, dichlofluanid, fluoromide, folpet or tolyfluanid; copper compounds such as Bordeaux mixture, copper hydroxide, copper oxychloride, copper sulfate, cuprous oxide, mancozeb or oxine-copper; nitrophenol derivatives such as dinocap or nitrothal-isopropyl; organo phosphorous derivatives such as edifenphos, iprobenphos, isoprothiolane, phosdiphen, pyrazophos or toclofos-methyl; and other compounds of diverse structures such as acibenzolar-S-methyl, anilazine, blasticidin-S, chinomethionat, chloroneb, chlorothalonil, cymoxanil, dichlone, diclomezine, dicloran, diethofencarb, dimethomorph, dithianon, etridiazole, famoxadone, fenamidone, fentin, ferimzone, fluazinam, flusulfamide, fenhexamid, fosetyl-aluminium, hymexazol, kasugamycin, methasulfocarb, pencycuron, phthalide, polyoxins, probenazole, propamocarb, pyroquilon, quinoxifen, quintozone, sulfur, triazoxide, tricyclazole, triforine, validamycin, (S)-5-methyl-2-methylthio-5-phenyl-3-phenyl-amino-3,5-dihydroimidazol-4-one (RPA 407213), 3,5-dichloro-N-(3-chloro-1-ethyl-1-methyl-2-oxopropyl)-4-methylbenzamide (RH-7281), N-allyl-4,5-dimethyl-2-trimethylsilylthiophene-3-carboxamide (MON 65500), 4-chloro-4-

cyano-N,N-dimethyl-5-p-tolylimidazole-1-sulfon-amide (IKF-916), N-(1-cyano-1,2-dimethyl-propyl)-2-(2,4-dichlorophenoxy)-propionamide (AC 382042), or iprovalicarb (SZX 722).

A preferred method of applying a compound of formula I, or an agrochemical composition which contains at least one of said compounds, is foliar application. The frequency of application and the rate of application will depend on the risk of infestation by the corresponding pathogen. However, the compounds of formula I can also penetrate the plant through the roots via the soil (systemic action) by drenching the locus of the plant with a liquid formulation, or by applying the compounds in solid form to the soil, e.g. in granular form (soil application). In crops of water rice such granulates can be applied to the flooded rice field. The compounds of formula I may also be applied to seeds (coating) by impregnating the seeds or tubers either with a liquid formulation of the fungicide or coating them with a solid formulation.

The formulation, i.e. the compositions containing the compound of formula I and, if desired, a solid or liquid adjuvant, are prepared in known manner, typically by intimately mixing and/or grinding the compound with extenders, e.g. solvents, solid carriers and, optionally, surface active compounds (surfactants).

The agrochemical formulations will usually contain from 0.1 to 99 % by weight, preferably from 0.1 to 95 % by weight, of the compound of formula I, 99.9 to 1 % by weight, preferably 99.8 to 5 % by weight, of a solid or liquid adjuvant, and from 0 to 25 % by weight, preferably from 0.1 to 25 % by weight, of a surfactant.

Advantageous rates of application are normally from 5 g to 2 kg of active ingredient (a.i.) per hectare (ha), preferably from 10 g to 1 kg a.i./ha, most preferably from 20 g to 600 g a.i./ha. When used as seed drenching agent, convenient dosages are from 10 mg to 1 g of active substance per kg of seeds.

Whereas it is preferred to formulate commercial products as concentrates, the end user will normally use dilute formulations.


The following non-limiting Examples illustrate the above-described invention in more detail. Temperatures are given in degrees Celsius. The following abbreviations are used:

m.p.= melting point; b.p.= boiling point. "NMR" means nuclear magnetic resonance spectrum. MS stands for mass spectrum. "%" is percent by weight, unless corresponding concentrations are indicated in other units.

Preparation Examples:

Example P 1: N-(2-biphenyl)-1-methyl-4-trifluoromethylpyrrol-3-carboxamide


a) 1-Methyl-4-trifluoromethylpyrrol-3-carboxylic acid



Sodium hydride (8.0g of a 75% dispersion in oil) is suspended at +5°C in a mixture of DMSO (300ml) and diethylether (100ml). A solution of ethyl 4,4,4-trifluorocrotonate (20g) and TOSMIC (23g) in DMSO (100ml) is added through a dropping funnel at such a rate that the temperature does not exceed 10°C. After stirring the reaction mixture for an additional hour at room temperature methyl iodide (15.6ml) is added with cooling. After 2 hours at room temperature the reaction mixture is poured onto crushed ice. Repeated extraction with ether, washing of the combined organic phases with brine and evaporation of the solvent under reduced pressure gives a product mixture in form of a light amber oil. The crude product mixture is heated at 60°C in a mixture of ethanol (100ml) and sodium hydroxide (50ml of a 30% aqueous solution). Washing of the solution with ether, acidifying of the aqueous phase with concentrated hydrochloric acid and filtering gives the 1-methyl-4-trifluoromethylpyrrol-3-carboxylic acid in form of a crystalline solid.

¹H-NMR(CDCl₃): 7.24(d,1H); 6.88(d,1H); 3.63 (s,3H).

b) N-(2-biphenyl)-1-methyl-4-trifluoromethylpyrrol-3-carboxamide




A solution of 1-methyl-4-trifluoromethylpyrrol-3-carboxylic acid (1.9g) and oxalyl chloride (0.9ml) in methylene chloride (20ml) is stirred at room temperature in the presence of a

catalytic amount of DMF. The solvent is evaporated under reduced pressure to give a crystalline solid. This solid is dissolved in methylene chloride (10ml) and added to a solution of 2-biphenylamine (1.7g) and triethylamine (4.2ml) in methylene chloride (20ml) at 0°C. The reaction mixture is stirred at room temperature for 2 hours. Evaporation of the solvent under reduced pressure, addition of water and filtering gives the N-(2-biphenyl)-1-methyl-4-trifluoromethylpyrrol-3-carboxamide.

¹H-NMR(CDCl₃): 8.37(d,1H); 7.60(s,br,1H); 7.48-7.14(m,8H); 7.03(d,1H); 6.88(d,1H); 3.63(s,3H).


Example P 2: N-(2-biphenyl)-1,5-dimethyl-4-trifluoromethylpyrrol-3-carboxamide

a) 1,5-Dimethyl-4-trifluoromethylpyrrol-3-carboxylic acid



A mixture of TOSMIC (20g), methyl iodide (18.5g) and N-benzyl triethylammonium chloride (1g) in methylene chloride (100ml) and aqueous sodium hydroxide (30ml of a 50% solution) are stirred at 5°C for 3 hours. A solution of ethyl 4,4,4-trifluorocrotonate (20g) in methylene chloride (50ml) is added through a dropping funnel at such a rate that the temperature does not exceed 25°C. After stirring the reaction mixture for an additional hour at room temperature excess methyl iodide is added with cooling. After 2 hours at room temperature the reaction mixture is poured onto crushed ice. Repeated extraction with ether, washing of the combined organic phases with brine and evaporation of the solvent gives the crude product mixture. The pure intermediate is obtained after chromatography on silicagel. Hydrolysis is achieved under the conditions mentioned under example 1 a) to give the 1,5-dimethyl-4-trifluoromethylpyrrol-3-carboxylic acid in form of a colorless crystalline solid. ¹H-NMR (d₆-DMSO): 12.04 (s,br,1H); 7.50(s,1H); 3.58(s,3H); 2.27 (s,3H).

b) N-(2-biphenyl)-1,5-dimethyl-4-trifluoromethylpyrrol-3-carboxamide



A solution of 1,5-dimethyl-4-trifluoromethylpyrrol-3-carboxylic acid (1.9g) and oxalyl chloride (0.9ml) in methylene chloride (20ml) is stirred at room temperature in the presence of a catalytic amount of DMF. The solvent is evaporated under reduced pressure to give a crystalline solid. This solid is dissolved in methylene chloride (10ml) and added to a solution of 2-biphenylamine (1.7g) and triethylamine (4.2ml) in methylene chloride (20ml) at 0°C. The reaction mixture is stirred at room temperature for 2 hours. Evaporation of the solvent under reduced pressure, addition of water and filtering gives the N-(2-biphenyl)-1,5-dimethyl-4-trifluoromethylpyrrol-3-carboxamide.

¹H-NMR(CDCl₃): 8.39(d,1H); 7.58(s,br,1H); 7.48-7.14(m,9H); 3.58 (s,3H); 2.27 (s,3H).

The following compounds are prepared in a similar way, using analogous methods.

Table 1: Compounds of formula I wherein R₁ is hydrogen, R₂ is methyl and A corresponds to one line in the Table A.

Table 2: Compounds of formula I wherein R₁ and R₂ are methyl and A corresponds to one line in the Table A.

Table 3: Compounds of formula I wherein R₁ is hydrogen, R₂ is ethyl and A corresponds to one line in the Table A.

Table 4: Compounds of formula I wherein R₁ is hydrogen, R₂ is methoxymethyl and A corresponds to one line in the Table A.

Table 5: Compounds of formula I wherein R₁ is hydrogen, R₂ is cyano and A corresponds to one line in the Table A.

Table 6: Compounds of formula I wherein R_1 is hydrogen, R_2 is methylsulfonyl and A corresponds to one line in the Table A.

Table 7: Compounds of formula I wherein R_1 is hydrogen, R_2 is phenylsulfonyl and A corresponds to one line in the Table A.




Table 8: Compounds of formula I wherein R_1 is hydrogen, R_2 is acetyl and A corresponds to one line in the Table A.

Table 9: Compounds of formula I wherein R_1 is hydrogen, R_2 is methoxyacetyl and A corresponds to one line in the Table A.


Table 10: Compounds of formula I wherein R_1 is hydrogen, R_2 is benzoyl and A corresponds to one line in the Table A.

Table 11: Compounds of formula I wherein R_1 is hydrogen, R_2 is 4-fluorobenzoyl and A corresponds to one line in the Table A.


Table A:

Comp. No.	A
01	
02	
03	


04




05




06




07




08




09




10




11




12


























13




14




15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	

27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	


38




39




40




41




42




43




44




45




46

























47



48



49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	

60	 <chem>CC(C)(C)CCc1cccc(C)c1</chem>
61	 <chem>CCC(C)c1cccc(C)c1</chem>
62	 <chem>CCC(C)(C)Cc1cccc(C)c1</chem>
63	 <chem>Clc1cc(Cl)ccc1Cc2cccc(C)c2</chem>
64	 <chem>c1ccc2ccccc2c1Cc3cccc(C)c3</chem>
65	 <chem>Cc1ccsc1Cc2ccc3ccccc3c2</chem>
66	 <chem>Cc1cccc2c1C=CC2</chem>
67	 <chem>CC1(C)C=CC2C1Cc3cccc(C)c3C2</chem>
68	 <chem>CC1(C)C=CC2C1Cc3cccc(C)c3C2C</chem>
69	 <chem>CC1(C)C=CC2C1Cc3cccc(C)c3C2C</chem>
70	 <chem>c1ccc(Oc2ccccc2)cc1C</chem>

71	
72	
73	
74	
75	
76	

Table 12 Physico-chemical data for selected compounds of the preceding tables 1 to 11

Comp. No.	physico-chemical data m.p.[°C] or ¹ H- NMR (CDCl ₃)
1.01	8.37(d,1H); 7.60(s,br,1H); 7.48-7.14(m,8H); 7.03(s,1H); 6.88(s,1H); 3.63 (s,3H); m.p. 100-102.
1.02	8.27(d,1H); 7.50(s,br,1H); 7.46-7.18(m,7H); 7.10(d,1H); 6.88(s,1H); 3.63(s,3H).
1.03	8.30(d,1H); 7.52(s,br,1H); 7.44-7.12(m,8H); 6.88(s,1H); 3.63(s,3H); m.p. 151-153.

1.04	7.85(d,1H); 7.74(s,br,1H); 7.44(dd, 2H); 7.30(s,1H); 7.28(s,1H); 7.13(dd,2H); 6.93(s,1H); 3.68(s,3H); m.p. 127-129.
1.05	7.84(d,1H); 7.74(s,br,1H); 7.49(s,4H); 7.30(d,1H); 7.28(d,1H); 6.96(d,1H); 3.68(s,3H); m.p. 142-145.
1.30	7.89(d,1H); 7.76-7.25(m,12H); 6.95(s,1H); 3.69(s,3H); m.p. 168-169.
1.35	m.p. 160-161.
1.38	8.25(d,1H); 7.50(s,br,1H); 7.45-7.08(m,7H); 6.68(s,1H); 3.65(s,3H); resin.
1.40	m.p. 122-123.
1.42	m.p. 142-143.
1.71	m.p. 63-65.
1.72	m.p. 112-114.
1.76	resin
2.02	m.p. 163-165.
2.03	resin
2.05	resin
2.35	m.p. 124-125.
2.71	resin
3.01	resin
3.02	resin
3.03	resin
3.71	resin
4.02	resin
4.03	resin
4.71	resin
5.02	resin
5.03	resin
5.71	resin
8.02	resin
8.03	resin
8.04	resin
8.05	resin
8.71	resin

Formulation Examples for compounds of formula I

Working procedures for preparing formulations of the compounds of formula I such as Emulsifiable concentrates, Solutions, Granulates, Dusts and Wettable powders are described in WO 97/33890.

Biological Examples: Fungicidal actions

Example B-1: Action against *Puccinia recondita* / wheat (Brownrust on wheat)

1 week old wheat plants cv. Arina are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application wheat plants are inoculated by spraying a spore suspension (1×10^5 uredospores/ml) on the test plants. After an incubation period of 2 days at 20° C and 95% r. h. plants are kept in a greenhouse for 8 days at 20° C and 60% r.h. The disease incidence is assessed 10 days after inoculation.

Compounds of Tables 1 to 11 show good activity in these tests (< 20% infestation).

Example B-2: Action against *Podosphaera leucotricha* / apple (Powdery mildew on apple)

5 week old apple seedlings cv. McIntosh are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. One day after application apple plants are inoculated by shaking plants infected with apple powdery mildew above the test plants. After an incubation period of 12 days at 22° C and 60% r. h. under a light regime of 14/10 h (light/dark) the disease incidence is assessed.

Compounds of Tables 1 to 11 show good activity in this test. The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit strong efficacy (< 20% infestation).

Example B-3: Action against *Venturia inaequalis* / apple (Scab on apple)

4 week old apple seedlings cv. McIntosh are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application apple plants are inoculated by spraying a spore suspension (4×10^5 conidia/ml) on the test plants. After an incubation period of 4 days at 21° C and 95% r. h. the plants are placed for 4 days at 21° C and 60% r. h. in a greenhouse. After another 4 day incubation period at 21° C and 95% r. h. the disease incidence is assessed.

Compounds of Tables 1 to 11 show good activity in this test. The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit strong efficacy (< 20% infestation).

Example B-4: Action against *Erysiphe graminis* / barley (Powdery mildew on barley)

1 week old barley plants cv. Express are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application barley plants are inoculated by shaking powdery mildew infected plants above the test plants. After an incubation period of 6 days at 20° C / 18° C (day/night) and 60% r. h. in a greenhouse the disease incidence is assessed.

Compounds of Tables 1 to 11 show good activity in this test. The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit strong efficacy (< 20% infestation).

Example B-5: Action against *Botrytis cinerea* / apple (Botrytis on apple fruits)

In an apple fruit cv. Golden Delicious 3 holes are drilled and each filled with 30 µl droplets of the formulated test compound (0.002% active ingredient). Two hours after application 50 µl of a spore suspension of *B. cinerea* (4×10^5 conidia/ml) are pipetted on the application sites. After an incubation period of 7 days at 22° C in a growth chamber the disease incidence is assessed.

Compounds of Tables 1 to 11 show good activity in this test. The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit very strong efficacy (< 10% infestation).

Example B-6: Action against *Botrytis cinerea* / grape (Botrytis on grapes)

5 week old grape seedlings cv. Gutedel are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. Two days after application grape plants are inoculated by spraying a spore suspension (1×10^6 conidia/ml) on the test plants. After an incubation period of 4 days at 21° C and 95% r. h. in a greenhouse the disease incidence is assessed.

Compounds of Tables 1 to 11 show good activity in this test . The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit very strong efficacy (< 10% infestation).

Example B-7: Action against *Botrytis cinerea* / tomato (Botrytis on tomatoes)

4 week old tomato plants cv. Roter Gnom are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. Two days after application tomato plants are inoculated by spraying a spore suspension (1×10^5 conidia/ml) on the test plants. After an incubation period of 4 days at 20° C and 95% r. h. in a growth chamber the disease incidence is assessed.

Compounds of Tables 1 to 11 show good activity in this test . The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit very strong efficacy (< 10% infestation).

Example B-8: Action against *Pyrenophora teres* / barley (Net blotch on barley)

1 week old barley plants cv. Express are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. Two days after application barley plants are inoculated by spraying a spore suspension (3×10^4 conidia/ml) on the test plants. After an incubation period of 2 days at 20° C and 95% r. h. plants are kept for 2 days at 20° C and 60% r.h. in a greenhouse. The disease incidence is assessed 4 days after inoculation.


Compounds of Tables 1 to 11 show good activity in this test . The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit strong efficacy (< 20% infestation).

Example B-9: Action against *Septoria nodorum* / wheat (Septoria leaf spot on wheat)

1 week old wheat plants cv. Arina are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application wheat plants are inoculated by spraying a spore suspension (5×10^5 conidia/ml) on the test plants. After an incubation period of 1 day at 20° C and 95% r. h. plants are kept for 10 days at 20° C and 60% r.h. in a greenhouse. The disease incidence is assessed 11 days after inoculation. Compounds of Tables 1 to 11 show good activity in this test . The compounds 1.01, 1.02, 1.03, 1.04, 1.05, 1.30, 1.35, 1.38, 1.39, 1.40, 1.42, 1.71, 1.72, 1.76, 2.02, 2.03, 2.05, 2.35, 2.71, 3.01, 3.02, 3.03, 3.71, 4.02, 4.03, 4.71, 5.02, 5.03, 5.71, 8.02, 8.03, 8.04 and 8.71 exhibit strong efficacy (< 20% infestation).

What is claimed is


1. A trifluoromethylpyrrolcarboxamide of formula I



wherein

R₁ is hydrogen, halogen, C₁₋₄haloalkyl or C₁₋₄alkyl,

R₂ is C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy-C₁₋₄alkyl, cyano, C₁₋₄alkylsulfonyl, phenylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, C₁₋₆alkylcarbonyl, benzoyl, or substituted phenylsulfonyl or benzoyl, and



wherein

R₃ is C₁₋₆alkyl, C₁₋₆haloalkyl, C₂₋₆alkenyl, C₂₋₆haloalkenyl, C₂₋₆alkinyl, C₁₋₆alkoxy, C₁₋₆haloalkoxy, C₂₋₆alkenyloxy, C₂₋₆haloalkenyloxy, C₂₋₆alkinyloxy, C₃₋₇cycloalkyl, C₁₋₄alkyl-C₃₋₇cycloalkyl, C₄₋₇cycloalkenyl, C₁₋₄alkyl-C₄₋₇cycloalkenyl, C₃₋₇cycloalkyloxy, C₁₋₄alkyl-C₃₋₇cycloalkyloxy, C₅₋₇cycloalkenyloxy, C₁₋₄alkyl-C₅₋₇cycloalkenyloxy, phenyl, naphthyl, phenoxy, naphthyloxy, or substituted phenyl or phenoxy wherein the substituents are one to three groups independently selected from halogen, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylthio, cyano, C₁₋₄alkoxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄haloalkyl, C₁₋₄haloalkoxy, methylenedioxy or difluoromethylenedioxy, or phenyl;

R₄ is hydrogen, halogen, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy or C₁₋₄haloalkoxy ; and


R₅, R₆ and R₇ independently of each other are C₁₋₆alkyl, C₃₋₇cycloalkyl or C₃₋₇cycloalkyl-C₁₋₄alkyl.

2. Compounds of formula I according to claim 1 wherein

R₁ is hydrogen or C₁₋₄alkyl,

R₂ is C₁₋₄alkyl, and

A is a group



wherein

R₃ is C₁₋₆alkyl, C₁₋₆haloalkyl, C₂₋₆alkenyl, C₂₋₆haloalkenyl, C₂₋₆alkinyl, C₁₋₆alkoxy, C₁₋₆haloalkoxy, C₂₋₆alkenyloxy, C₂₋₆haloalkenyloxy, C₂₋₆alkinyloxy, C₃₋₇cycloalkyl, C₁₋₄alkyl-C₃₋₇cycloalkyl, C₄₋₇cycloalkenyl, C₁₋₄alkyl-C₄₋₇cycloalkenyl, C₃₋₆cycloalkyloxy, C₁₋₄alkyl-C₃₋₇cycloalkyloxy, C₅₋₇cycloalkenyloxy, C₁₋₄alkyl-C₅₋₇cycloalkenyloxy, phenyl, naphthyl, phenyl substituted by one to three groups independently selected from halogen, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylthio, cyano, C₁₋₄alkoxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄haloalkyl, C₁₋₄haloalkoxy, methylenedioxy or difluoromethylenedioxy, or phenyl;


R₄ is hydrogen, halogen or C₁₋₄alkyl ;and

R₅, R₆ and R₇ independently of each other are C₁₋₆alkyl, C₃₋₇cycloalkyl or C₃₋₇cycloalkyl-C₁₋₄alkyl.

3. A compound according to claims 1 or 2 wherein R₁ is hydrogen or methyl.
4. A compound according to claims 1 or 2 wherein R₂ is methyl.
5. A compound according to claims 1 or 2 wherein R₃ is phenyl or phenyl substituted with halogen.
6. A compound according to claims 1 or 2 wherein R₁ is hydrogen or methyl, R₂ is methyl, and R₃ is phenyl or phenyl substituted with halogen.
7. A compound according to claim 1 selected from the group comprising
 N-(2-biphenyl)-1-methyl-4-trifluoromethylpyrrol-3-carboxamide,
 N-(4'-chloro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide,
 N-(4'-fluoro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide,
 N-[2-(4-fluorophenyl)-3-thienyl]-4-trifluoromethylpyrrol-3-carboxamide,


N-[2-(4-chlorophenyl)-3-thienyl]-4-trifluoromethylpyrrol-3-carboxamide,
 N-(3',4'-difluoro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide,
 N-(3'-trifluoromethyl-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide, and
 N-(4'-chloro-3'-fluoro-2-biphenyl)-4-trifluoromethylpyrrol-3-carboxamide.

8. A process for the preparation of compounds of formula I which comprises reacting the starting materials according to the scheme



wherein A, R_1 and R_2 are as defined for formula I in claim 1, X and Y are leaving groups, alkyl designates a lower alkyl moiety, and aryl stands for phenyl or tolyl.

9. A process for the preparation of compounds of formula I which comprises reacting the starting materials according to the scheme



wherein A, R_1 and R_2 are as defined for formula I in claim 1, X and Y are leaving groups, and alkyl designates a lower alkyl moiety, optionally with the utilization of a protective group during the amidation step.

10. A composition for controlling microorganisms and preventing attack and infestation of plants therewith, wherein the active ingredient is a compound as claimed in claim 1 together with a suitable carrier.
11. Use of a compound of formula I according to claim 1 for protecting plants against infestation by phytopathogenic microorganisms.
12. A method of controlling or preventing infestation of cultivated plants by phytopathogenic microorganisms by application of a compound of formula I as claimed in claim 1 to plants, to parts thereof or to the locus thereof.

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/EP 99/05837

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D207/40 C07D409/12 C07D409/14 A01N43/36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 08148 A (BASF A.-G., GERMANY) 6 March 1997 (1997-03-06) abstract; claims page 23; examples 3.9-3.11; table 3 ---	1,10-12
Y	EP 0 737 682 A (MITSUI TOATSU CHEMICALS, INCORPORATED, JAPAN) 16 October 1996 (1996-10-16) abstract; claims page 27; examples 1.1,1.2,1.4,1.6; table 5 ---	1,10-12
Y	EP 0 566 138 A (HODOGAYA CHEMICAL CO., LTD., JAPAN) 20 October 1993 (1993-10-20) abstract; claims page 12 -page 13; examples 77,79; table 1 --- -/--	1,10-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 February 2000

Date of mailing of the international search report

10/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Paisdor, B

INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/EP 99/05837

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>EP 0 545 099 A (BASF AG) 9 June 1993 (1993-06-09) cited in the application abstract; claims page 23 page 34 -page 36; table 10 page 6; examples 1.75-1.77; table -----</p>	1, 10-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05837

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9708148 A	06-03-1997	DE 19531813 A	06-03-1997
		AU 6928596 A	19-03-1997
		EP 0847388 A	17-06-1998
		JP 11511449 T	05-10-1999
		US 5998450 A	07-12-1999
EP 0737682 A	16-10-1996	CA 2173788 A	12-10-1996
		CN 1146993 A	09-04-1997
		JP 9235282 A	09-09-1997
		US 5747518 A	05-05-1998
EP 0566138 A	20-10-1993	DE 69322118 D	24-12-1998
		DE 69322118 T	20-05-1999
		JP 6025199 A	01-02-1994
EP 0545099 A	09-06-1993	AT 149487 T	15-03-1997
		AU 656243 B	27-01-1995
		AU 2855492 A	27-05-1993
		CA 2081935 A	23-05-1993
		DE 59208113 D	10-04-1997
		DK 545099 T	24-03-1997
		ES 2098421 T	01-05-1997
		GR 3023336 T	29-08-1997
		HU 213622 B	28-08-1997
		IL 103614 A	24-09-1998
		JP 5221994 A	31-08-1993
		NZ 245194 A	27-02-1996
		PL 296677 A	18-10-1993
		SK 344892 A	08-03-1995
		US 5480897 A	02-01-1996
		US 5556988 A	17-09-1996
		US 5589493 A	31-12-1996
		US 5330995 A	19-07-1994
		ZA 9208977 A	19-05-1994